Black Holes with Anisotropic Fluid

Inyong Cho[1,*] and Hyeong-Chan Kim[2,†]

[1]*School of Liberal Arts, Seoul National University of Science and Technology, Seoul 01811, Korea*
[2]*School of Liberal Arts and Sciences, Korea National University of Transportation, Chungju 27469, Korea*

We study a spherically symmetric spacetime made of an anisotropic perfect fluid for which the sum of the radial pressure and the energy density vanishes. For a given equation-of-state parameter $w_2 = p_2/\rho$, we find exact solutions of the Einstein's equation which are parametrized by two independent parameters M and Λ which play the role of the mass and the cosmological constant in the isotropic case. Then, we classify the solution into six types based on the behavior of the metric function. Depending on the parameters, the solution can have event horizons and a cosmological horizon.

PACS numbers: 04.70.Bw, 04.20.Jb
Keywords: black hole, exact solution, general relativity

arXiv:1703.01103v1 [gr-qc] 3 Mar 2017

I. INTRODUCTION

A comprehensive collection of static solutions of Einstein's field equation with spherical symmetry can be found in Stephani *et. al.* [1], Delgaty and Lake [2], and Semiz [3]. Most of them focused on the isotropic fluids because astrophysical observations support isotropy. The perfect Pascalian-fluid (isotropic-fluid) assumption is supported by solid observational and theoretical grounds. However, an increasing amount of theoretical evidence strongly suggests that, for certain density ranges, a variety of very interesting physical phenomena giving rise to local anisotropy may take place (see [4] and references therein). As investigated by Ruderman [5], highly compact astrophysical objects having core density beyond the nuclear density ($\sim 10^{15}\mathrm{g/cm}^3$) can have pressure anisotropy, i.e., the pressure inside these compact objects can be decomposed into two parts, the radial pressure p_1 and the transverse pressure p_2 in two perpendicular directions to p_1. There are various underlying reasons of this anisotropic nature, for example, the existence of solid core in the presence of type 3A superfluid [6]. The local anisotropy in self-gravitating systems were studied [4]. The pressure anisotropy affects the physical properties such as the stability and the structure of stellar matter [7]. Recently, Bhar [8, 9] studied a new model of an anisotropic superdense star which admits conformal motions in the presence of a quintessence field which is characterized by the equation-of-state parameter w with $-1 < w < -1/3$. The self-gravitating charged anisotropic fluid with barotropic equation of state were also considered [10–12]. The Einstein's field equation was solved by assuming some specific mass function in Refs. [9, 13, 14].

Following these recent trend, in this work, we consider the anisotropic fluid to find exact solutions of the Einstein's field equation. For simplicity, we consider the static and spherical symmetric configurations. Then, the metric ansatz becomes

$$ds^2 = -f(r)dt^2 + g(r)dr^2 + r^2 d\theta^2 + r^2 \sin^2\theta\, d\phi^2. \tag{1}$$

The simplifying assumptions mostly used for matter are the vacuum, the electromagnetic field, and the perfect fluid. For example, the vacuum with the ansatz (1) gives uniquely the Schwarzschild metric, the simplest and best-known black-hole solution. The stress tensor for an anisotropic perfect fluid compatible with the spherical symmetry is

$$T_{\mu\nu} = (\rho + p_2)u_\mu u_\nu + (p_1 - p_2)x_\mu x_\nu + p_2 g_{\mu\nu}, \tag{2}$$

where ρ is the energy density measured by a comoving observer with the fluid, and u^μ and x^μ are its four-velocity and a spacelike unit vector orthogonal to u^μ and angular directions, respectively. The use of this $T_{\mu\nu}$ together with ansatz (1) can describe the interior of static, spherically symmetric, and extremely high-density stars, for example. In addition to these symmetry, we need to introduce the equation of state, which is the relation between p_i and ρ. In cosmology, one usually assumes the barotropic condition for the equation of state,

$$p_i = w_i \rho, \tag{3}$$

*Electronic address: iycho@seoultech.ac.kr
†Electronic address: hckim@ut.ac.kr

with $w_i = 0$ describing the dust, $w_i = 1/3$ the radiation, $w_i < -1/3$ the dark energy, and $w_i < -1$ the phantom energy.

Some general requirements for $T_{\mu\nu}$ are proposed, collectively known as *energy conditions*. For example, the weak energy condition states that the energy density should be nonnegative to every observer. Alternatively, one might impose strong or dominant energy conditions. For the anisotropic fluid, the energy conditions take the following forms: the weak energy condition, $\rho \geq 0$, $\rho + p_i \geq 0$, the strong energy condition, $\rho + p_i \geq 0$, $\rho + \sum_i p_i \geq 0$, the dominant energy condition, $\rho \geq |p_i|$, and the null energy condition, $\rho + p_i \geq 0$. When $g(r) < 0$ and $f(r) < 0$, the coordinate r plays the role of time. Then, $-p_1$ and $-\rho$ play the roles of the energy density and the pressure along the spatial t direction, respectively. Upon this swich of the roles, the energy conditions mentioned above do not change and the energy density and the pressure are continuous across the horizon when $w_1 = -1$. We are interested in this case at the present work. For other cases, $w_1 \neq -1$, the pressure must be discontinuous at the horizon r_H unless $\rho(r_H) = 0$. In Ref. [2], various tests of acceptability for the isotropic fluid, such as the positivity of energy density and pressure, the regularity at the origin, the subluminal sound speed, etc. were applied to vast of candidates for fluid solutions.

In this work, we consider the case of $w_1 = -1$ for which there exist exact analytic solutions. We classify the solutions according to the value of w_2. In Sec. II, we derive the Tolman-Oppenheimer-Volkoff equation for the anisotropic fluid. In Sec. III, we obtain the exact solution for the case of $w_1 = -1$. In Sec. IV, we classify the solution into six types. In Sec. V, we discuss the black-hole types. In Sec. VI, we study the stability of the solutions. We summarize the results in Sec. VII.

II. FIELD EQUATION

With the metric (1) and the energy-momentum tensor (2), the Einstein's equation becomes

$$G^0_0 = -\frac{1}{r^2} + \frac{1}{r^2 g} - \frac{g'}{rg^2} = -8\pi\rho(r), \tag{4}$$

$$G^1_1 = -\frac{1}{r^2} + \frac{1}{r^2 g} + \frac{f'}{rfg} = 8\pi p_1(r), \tag{5}$$

$$G^2_2 = \frac{f'}{2rfg} - \frac{f'^2}{4f^2 g} - \frac{g'}{2rg^2} - \frac{f'g'}{4fg^2} + \frac{f''}{2fg} = 8\pi p_2(r). \tag{6}$$

The first equation (4) can be formally integrated to give

$$g(r) = \frac{1}{1 - 2m(r)/r}, \tag{7}$$

where the mass function $m(r)$ is defined by

$$m(r) = 4\pi \int^r r'^2 \rho(r') dr'. \tag{8}$$

Here, the integration constant is absorbed into the definition of $m(r)$. Using Eqs. (4), (5) and (6), or equivalently the conservation law $\nabla^\mu T_{\mu\nu} = 0$, we obtain the hydrostatic equation for anisotropic fluids

$$\frac{1}{2}\frac{f'}{f}(\rho + p_1) + p_1' + \frac{2(p_1 - p_2)}{r} = 0. \tag{9}$$

Using Eq. (3), this equation can be integrated to give

$$\rho(r) = \rho_0 f^{-(1+w_1)/2w_1} r^{-2(w_1 - w_2)/w_1} \equiv \frac{m'}{4\pi r^2}. \tag{10}$$

Integrating Eq. (5) by using $p_1(r) = w_1\rho(r)$ and Eq. (4), we write the metric functions $f(r)$ as a function of $m(r)$,

$$f(r) = \frac{[g(r)]^{w_1}}{r^{w_1+1}} \exp\left[(1+w_1)\int^r \frac{g(r')}{r'} dr'\right] = \frac{[r - 2m(r)]^{-w_1}}{r} \exp\left[(1+w_1)\int^r \frac{1}{r' - 2m(r')} dr'\right]. \tag{11}$$

Plugging Eqs. (7) and (11) into Eq. (6), and using $p_1 = w_1\rho$ and $p_2 = w_2\rho$, we get the Tolman-Oppenheimer-Volkoff (TOV) equation [15] for the anisotropic fluid,

$$\frac{m''}{m'} + \frac{1 + w_1}{r - 2m}\left(m' + \frac{1}{2w_1}\right) - \frac{1 + w_1 + 4w_2}{2w_1 r} = 0. \tag{12}$$

For the isotropic case, it becomes the usual TOV equation written in terms of m.

III. ANALYTIC SOLUTIONS

The purpose of this work is to find analytic solutions of the TOV equation (12). For the isotropic case with $w_1 = w_2 = -1/3$, analytic solutions including the black hole in closed spacetime were found in Ref. [16]. Another exact analytic solution, we are interested in this work, can be found when

$$w_1 = -1. \tag{13}$$

When ρ plays the role of an energy density, the energy conditions restrict the matter kinds to physically allowed ones. Among the conditions, the positivity of energy density appears to be crucial. In addition to it, all the energy conditions require $w_2 \geq -1$. Specifically, the dominant energy condition requires $w_2 \leq 1$ and the strong energy condition requires $w_2 \geq 0$. Therefore, when $0 \leq w_2 \leq 1$, all the energy conditions will be satisfied.

With Eq. (13), the TOV equation (12) takes integrable form, $m''/m' + 2w_2/r = 0$. The solution is given by

$$m(r) = \begin{cases} \frac{\Lambda}{2} r^{1-2w_2} + M & (w_2 \neq \frac{1}{2}) \\ \frac{\Lambda}{2} \log\left|\frac{r}{\Lambda}\right| + M & (w_2 = \frac{1}{2}) \end{cases}. \tag{14}$$

This gives the density and the radial pressure as

$$\rho(r) = -p_1(r) = \begin{cases} \frac{(1-2w_2)\Lambda}{8\pi r^{2w_2+2}} & (w_2 \neq \frac{1}{2}) \\ \frac{\Lambda}{8\pi r^3} & (w_2 = \frac{1}{2}) \end{cases}. \tag{15}$$

The energy density and the pressure are singular at the origin and at the infinity when $w_2 > -1$ and $w_2 < -1, \neq 1/2$, respectively. Now, $\rho(r)$ is positive definite when (i) $(w_2 - 1/2)\Lambda < 0$ and (ii) $\Lambda > 0$ and $w_2 = 1/2$. The metric functions in Eqs. (7) and (11) become

$$g(r)^{-1} = f(r) = \begin{cases} 1 - \frac{2M}{r} - \frac{\Lambda}{r^{2w_2}}, & (w_2 \neq \frac{1}{2}) \\ 1 - \frac{2M}{r}\left(1 + \frac{\Lambda}{2M} \log\left|\frac{r}{\Lambda}\right|\right), & (w_2 = \frac{1}{2}) \end{cases}. \tag{16}$$

When $\Lambda = 0$ the energy density vanishes and the metric becomes nothing but the Schwarzschild spacetime. Because we are interested in the solutions with matter, we restrict our interests to the case with $\Lambda \neq 0$. For the isotropic fluid with $w_1 = w_2 = -1$, M and Λ represent the mass of the (anti)-de Sitter black hole and the cosmological constant, respectively.

Now, let us analyze the curvature singularities. The scalar curvature,

$$R = \frac{2\Lambda(w_2 - 1)(2w_2 - 1)}{r^{2(w_2+1)}},$$

is singular at the origin and at the infinity when $-1 < w_2 \neq 1, 1/2$ and $w_2 < -1$, respectively. Therefore, it is regular everywhere only when $w_2 = -1, 1$ and $1/2$. The Krestchmann invariant is given by

$$R_{abcd}R^{abcd} = \frac{48M^2}{r^6} + \frac{16\Lambda M(w_2 + 1)(2w_2 + 1)}{r^{2w_2+5}} + \frac{4\Lambda^2\left(4w_2^4 + 4w_2^3 + 5w_2^2 + 1\right)}{r^{4w_2+4}}.$$

Note that the numerator of the last term is positive definite. Therefore, the term is singular at the origin or at the infinity when $w_2 > -1$ or $w_2 < -1$, respectively. When $w_2 = -1$, the first term becomes singular at the origin unless $M = 0$. For $\Lambda \neq 0$, the Krestchmann invariant is regular everywhere only when $M = 0$ and $w_2 = -1$, of which solution is nothing but the (anti)-de Sitter spacetime. In summary, the spacetime is singular at infinity when $w_2 < -1$ and singular at the origin when $M \neq 0$. The singularities can be naked or be covered by (cosmological) event horizons depending on the natures of the spacetime.

IV. CLASSIFICATION OF SOLUTIONS

We classify the solution in Eq. (16) into six types in accordance with the behaviors of the function $f(r)$ as following.
(I) Schwarzschild: $f(r)$ changes the signature from negative to positive with r.
(II) de Sitter: $f(r)$ changes the signature from positive to negative.
(III) Schwarzschild-de Sitter: $f(r)$ changes the signature from negative to positive and then to negative again.



FIG. 1: Typical form of $f(r)$ for type I-VI.

(IV) Two horizon black hole: $f(r)$ changes the signature from positive to negative and then to positive again.
(V) Static naked: $f(r)$ is positive definite.
(VI) Time-dependent singular: $f(r)$ is negative definite.

Before we discuss the details of each solution, we would like to discuss the positivity of the energy density when there exists an event horizon in the solution. As $w_1 < 0$, the signature of ρ is different from that of p_1. Because t and r coordinates exchange their roles of time and space, $-p_1$ plays the role of the energy density in the region of $f(r) < 0$. Therefore, in the presence of a horizon, the energy density is positive definite over the whole space when $w_1 < 0$ and $\rho > 0$. For the present case that we consider, the condition $\rho > 0$ in Eq. (15) requires $w_2 > 1/2$ for $\Lambda < 0$, or $w_2 \leq 1/2$ for $\Lambda > 0$.

The full classification of the solutions with respect to the schemes above is given in Fig. 2. The Roman numbers I-VI represent type of the given solution corresponding to the given values of (w_2, M, Λ). The character X represents that there are no region having a positive definite energy density in the solution.



(a) $w_2 < 0$ (b) $w_2 = 0$ (c) $0 < w_2 < \frac{1}{2}$ (d) $w_2 = \frac{1}{2}$ (e) $w_2 > \frac{1}{2}$

FIG. 2: The classification of the solutions. The horizontal and the vertical axes denote M and Λ, respectively. The curves in (a), (c), and (e) are given by the formula $(2w_2 - 1)M/w_2 = [(1 - 2w_2)\Lambda]^{1/2w_2}$. The straight lines in (b) and (d) represent $\Lambda = 1$ and $\Lambda = 2M$, respectively. Each border line in the figure belongs to the part having the same text color.

To illustrate the figure, let us consider the case of $w_2 > 1/2$ in detail as an example. ρ is positive definite when Λ is negative. Therefore, the positive Λ region is not physical. Around the origin, $r \sim 0$, the behavior of the function $f(r)$ is governed by the last term containing Λ. On the other hand, $f(r) \to 1$ asymptotically as $r \to \infty$. If Λ is negative, $f(r)$ is a decreasing function with $f(0) = \infty$. Its derivative is

$$f'(r) = \frac{2M}{r^2} + \frac{2w_2\Lambda}{r^{2w_2+1}}.$$

When M is positive, $f'(r)$ vanishes at $r_m = (-w_2\Lambda/M)^{1/(2w_2-1)}$ and the value of $f(r)$ at r_m is

$$f(r_m) = 1 - \frac{2M}{r_m} - \frac{\Lambda}{r_m^{2w_2}} = 1 - \frac{2w_2 - 1}{w_2}\frac{M}{r_m} = 1 - \left\{\frac{(2w_2 - 1)M}{w_2}[(1 - 2w_2)\Lambda]^{-1/(2w_2)}\right\}^{2w_2/(2w_2-1)}.$$

The minimum value $f(r_m)$ is positive when $(2w_2-1)M/w_2 < [(1-2w_2)\Lambda]^{1/(2w_2)}$. For this case, the metric function is positive definite for all r and the singularity at the origin is naked. Now, the metric belongs to type V. The case of $M<0$ also belongs to this type. On the other hand, when $f(r_m)<0$, the spacetime has two horizons at the radius $r_\pm$ satisfying $f(r_\pm)=0$. Therefore, the spacetime belongs to type IV.

If Λ is positive, the energy density (15) is negative definite. Therefore, the metrics are unphysical. $f(r)$ increases from $f(0)=-\infty$. $f(r)$ may have a maximum at a point r_M, or may not. The maximum value must be larger than unity because $f(r)\to 1$ asymptotically and the equation $f'(r)=0$ can have at most one root. Therefore, $f(r)$ vanishes only at one place (r_H) for any value of M. This implies that the solution belongs to type I X.

The solutions corresponding to a black hole are type I, III, and IV. The type II solution is similar to that of the de Sitter spacetime having a cosmological horizon. On the other hand, the type V solution possesses a naked singularity. The type VI solution corresponds to a cosmological solution undergoing anisotropic expansion/contraction which is bounded by a cosmological horizon. The singularity at $r=0$ corresponds to the future/past infinity in this case.

V. BLACK-HOLE SOLUTIONS

Out of the six types, only the types I, III, and IV correspond to the black-hole spacetime. Because the black-hole solutions are most interesting, we focus on these three classes in this section.

A. Type I

This type of solutions exists only when $0\le w_2\le 1/2$ and $\Lambda>0$. For this range, the energy density is positive definite, $\rho>0$, everywhere. The parameter M is positive for $w_2<1/2$ and is negative for $w_2=1/2$. The geometry takes the form of a modified Schwarzschild spacetime. The functional form of $f(r)$ at $r\simeq 0$ is governed by the $-2M/r$ term, which implies that $r=0$ is a singularity. The geometry is asymptotically flat because $f(r)$ approaches a constant value as $r\to\infty$. There exists an event horizon $r_{\rm BH}$ satisfying $f(r_{\rm BH})=0$.

If $w_2=0$, the solution takes the form of a modified Schwarzschild form with a solid angle deficit. After introducing a new coordinates $y=r/\alpha$ and $\tau=\alpha t$, where $\alpha=\sqrt{1-\Lambda}$, the metric becomes

$$ds^2=-\left(1-\frac{2M'}{y}\right)d\tau^2+\frac{dy^2}{1-2M'/y}+(\alpha y)^2(d\theta^2+\sin^2\theta d\phi^2),$$

where $M'=M/\alpha^3$. The horizon is at $r=2M/\alpha^2$. The mass function increases with r as $m(r)=M+\Lambda r/2$. At the horizon, it is $m(r_{\rm BH})=M/\alpha^2$. The solution mimics that of the global monopole [17].

B. Type III

The type III spacetime exists only for $w_2<0$ with Λ and M being positive definite. The geometry takes the form of the modified Schwarzschild-de-Sitter solution. The functional form of $f(r)$ at $r\sim 0$ is governed by the $-2M/r$ term, which implies that $r=0$ is a singularity. Asymptotically, $f(r)\to-\Lambda r^{2|w_2|}$ goes to negative infinity. $f(r)$ has a maximum

$$f(r_m)=1-\frac{2|w_2|+1}{|w_2|}\frac{M}{r_m},\qquad r_m=\left|\frac{M}{\Lambda w_2}\right|^{1/(2|w_2|+1)}.$$

For this type, the maximum value $f(r_m)$ is positive. There are two horizons satisfying $f(r)=0$, the one is the black-hole horizon at $r_{\rm BH}\le r_m$ and the other is the cosmological horizon at $r_H\ge r_m$. The cosmological horizon resembles that of the de-Sitter spacetime. If $f(r_m)=0$, the cosmological horizon coincides with the black-hole horizon. When $f(r_m)\le 0$, the solution belongs to Type VI.

When $w_2>-1$, $\rho(r)$ diverges at $r=0$ then monotonically decreases to zero at infinity. On the other hand, when $w_2<-1$, it monotonically increases from zero and diverges at infinity. When $w_2=-1$, the geometry becomes the Schwarzschild-de-Sitter spacetime with $f(r)=1-2M/r-\Lambda r^2$.

C. Type IV

The type IV spacetime exists only for $w_2>0$. The values of M and Λ can take both signatures depending on the value of w_2. The geometry of a type IV solution takes the form of the Reissner-Nordström solution in the sense that

it has an inner horizon in addition to the outer black-hole horizon. At the origin, $f(r)$ diverges and the spacetime is asymptotically flat, i.e., $f(r) \to 1$. $f(r)$ has a minimum $f(r_m)$ at $r = r_m$. When $w_2 \neq 1/2$,

$$f(r_m) = 1 - \frac{2w_2 - 1}{w_2}\frac{M}{r_m} \leq 0, \qquad r_m = \left|\frac{M}{-\Lambda w_2}\right|^{1/(1-2w_2)}.$$

The outer horizon is located at $r_{\rm BH} \geq r_m$. When $|\Lambda/M| = w_2^{-1}\left[(2w_2-1)M/w_2\right]^{2w_2-1}$, the two horizons merge to form an extremal black hole. If the ratio $|\Lambda/M|$ is smaller than this value, the singularity becomes naked and the solution becomes type V. When $w_2 = 1$, the solution becomes the exact Reissner-Nordstorm metric with charge $Q^2 = -\Lambda$.

VI. STABILITY

Let us check the stability of the spacetime with anisotropic fluid by introducing linear spherical scalar perturbations. Let us write the metric in the form,

$$ds^2 = -e^{\nu}dt^2 + e^{\lambda}dr^2 + e^{\mu}d\Omega_2^2. \tag{17}$$

The general energy-momentum tensor is given by Eq. (2) with $p_i = w_i\rho$, where the velocity and the radial four vectors are given by

$$u^\mu = [e^{-\nu/2}\sqrt{1+v^2}, e^{-\lambda/2}v, 0, 0], \qquad x^\nu = [e^{-\nu/2}v, e^{-\lambda/2}\sqrt{1+v^2}, 0, 0], \tag{18}$$

where v corresponds to a normalized radial velocity and $x^a u_a = 0$. Now, the components of the stress tensor become

$$T_1^0 = (1+w_1)\rho e^{(\lambda-\nu)/2}v\sqrt{1+v^2}, \quad T_0^0 = -\rho\left[1+(1+w_1)v^2\right], \quad T_1^1 = \rho[w_1+(1+w_1)v^2], \quad T_2^2 = \rho w_2. \tag{19}$$

Now, we introduce the perturbations for the metric on a given background solution $\nu_0(r)$, $\lambda_0(r)$, and $\mu_0(r)$ as

$$e^\nu = e^{\nu_0(r)}(1+\delta\nu), \qquad e^\lambda = e^{\lambda_0(r)}(1+\delta\lambda), \qquad e^\mu = e^{\mu_0(r)}(1+\delta\mu), \qquad \rho = \rho_0 + \delta\rho. \tag{20}$$

The linearized coordinate transformations of t and r are given by

$$t = \tilde{t} + \delta t(\tilde{t}, \tilde{r}), \qquad r = \tilde{r} + \delta r(\tilde{t}, \tilde{r}). \tag{21}$$

Under this transformation, the metric (17) can be written as

$$\begin{aligned} ds^2 &= -e^{\nu_0(r)}(1+\delta\nu)dt^2 + e^{\lambda_0(r)}(1+\delta\lambda)dr^2 + e^{\mu_0(r)}(1+\delta\mu)d\Omega^2 \\ &= -e^{\nu_0(\tilde{r})}[1+\delta\nu+\nu_0'\delta r + \dot{\nu}_0\delta t + 2\dot{\delta t}]d\tilde{t}^2 + 2[e^{\lambda_0}\dot{\delta r} - e^{\nu_0}\delta t']d\tilde{t}d\tilde{r} \\ &\quad + e^{\lambda_0(\tilde{r})}[1+\delta\lambda+\lambda_0'\delta r + \dot{\lambda}_0\delta t + 2\delta r']d\tilde{r}^2 + e^{\mu_0(\tilde{r})}\left[1+\delta\mu+\mu_0'\delta r + \dot{\mu}_0\delta t\right]d\Omega^2, \end{aligned} \tag{22}$$

where the overdot and the prime denote the derivatives with respect to $\tilde{t}$ and $\tilde{r}$, respectively. In Eq. (22), we can impose $\dot{\nu}_0 = 0$, $\dot{\lambda}_0 = 0$, and $\dot{\mu}_0 = 0$.

In this work, we choose the gauge choice for δr and δt so that

$$\tilde{g}_{0i} = 0 \quad \Rightarrow \quad e^{\lambda_0}\dot{\delta r} = e^{\nu_0}\delta t', \qquad \delta\tilde{\mu} \equiv \delta\mu + \mu_0'\delta r = 0. \tag{23}$$

Omitting tilde in $\tilde{r}$ and $\tilde{t}$, the metric (22) becomes

$$ds^2 = -e^{\nu_0}[1+\delta\nu(t,r)]dt^2 + e^{\lambda_0}[1+\delta\lambda(t,r)]dr^2 + e^{\mu_0}d\Omega^2. \tag{24}$$

The δG_1^0 part of the Einstein's equation based on the metric (24), introducing $e^{\nu_0} = f(r) = e^{-\lambda_0}$, determines v to be

$$-\frac{\mu_0'\dot{\delta\lambda}}{2f(r)} = 8\pi(1+w_1)\frac{\rho_0 v}{f} \quad \Rightarrow \quad v = -\frac{\mu_0'\,\dot{\delta\lambda}}{16\pi(1+w_1)\rho_0}. \tag{25}$$

We may use this equation as a definition of v. The δG_0^0 part,

$$-\frac{f(r)}{4}\left[2\mu_0'\delta\lambda' + \left(4\mu_0'' + 3\mu_0'^2\right)\delta\lambda\right] = -8\pi\delta\rho, \tag{26}$$

provides the definition of $\delta\rho$. The δG^1_1 part becomes

$$\frac{f(r)}{4}\left(-\mu_0'^2\delta\lambda + 2\mu_0'\delta\nu'\right) = 8\pi w_1\delta\rho. \tag{27}$$

Combining Eq. (26) and Eq. (27), we write $\delta\nu'$ using $\delta\lambda$:

$$\delta\nu' = w_1\delta\lambda' + \Big(\frac{1+3w_1}{2} + 2w_1\frac{\mu_0''}{\mu_0'^2}\Big)\mu_0'\,\delta\lambda. \tag{28}$$

Finally, the δG^2_2 part becomes

$$-\frac{\ddot{\delta\lambda}}{2f(r)} + \frac{f(r)}{4}\left[2\delta\nu'' + \mu_0'(\delta\nu' - \delta\lambda') - (2\mu_0'' + \mu_0'^2)\delta\lambda\right] = 8\pi w_2\delta\rho. \tag{29}$$

Using $\mu_0(r) = 2\log r$ after plugging Eqs. (28) and (27) into Eq. (29), we get a differential equation for the perturbation $\delta\lambda$ as

$$\ddot{\delta\lambda} = f^2(r)\left[w_1\delta\lambda'' + \frac{w_1 - w_2}{r}\delta\lambda' - \frac{2w_2}{r^2}\delta\lambda\right]. \tag{30}$$

Introducing coordinate z as $dz = f^{-1}dr$ and setting $\delta\lambda = e^{-i\omega t}g_1(r)$, the perturbation equation becomes

$$-\frac{d^2g_1}{dz^2} + \frac{f'}{w_1}\frac{dg_1}{dz} + V(z)g_1 = \frac{\omega^2}{w_1}g_1, \qquad V(z) \equiv \frac{2w_2}{w_1}\frac{f^2(r)}{r^2}. \tag{31}$$

An interesting point here is that the pressure orthogonal to the radial direction enters into the potential part only. For large ω^2, the equation takes the form, $d^2g_1/dz^2 \approx (-w_1^{-1})\omega^2 g_1$. Therefore, the system has possibility for being stable only when $w_1 \geq 0$. For the present system, $w_1 = -1$ implies that the system must be unstable under the radial perturbations.

VII. SUMMARY

We have studied static geometries with an anisotropic perfect fluid based on Einstein's theory of gravity. Because of the spherical symmetry, the angular pressures should be isotropic. However, it can be different from the radial pressure without breaking the spherical symmetry. The Tolman-Oppenheimer-Volkoff (TOV) equation is obtained for linear equation of state of the system. When $p_1 = -\rho$, the TOV equation takes simple form, $m''/m' + 2w_2/r = 0$, of which solution allows various classes with $p_2 = w_2\rho$.

Formally, by using the areal radial coordinate, the solution takes similar form with that of the Schwarzschild-de Sitter-like spacetime,

$$g_{rr}^{-1} = g_{tt} = 1 - \frac{2M}{r} - \frac{\Lambda}{r^{2w_2}}.$$

For the given equation-of-state parameter w_2, the spacetime geometry is described by two different parameters M and Λ, representing the mass and the cosmological constant for the isotropic case. When $M \neq 0$, the solution has a curvature singularity at $r = 0$. It additionally has a singularity at infinity when $w_2 < -1$ and $\Lambda \neq 0$. On the basis of the behavior of the metric function, we classified the solutions into six types, including black holes (types I, III, and IV), de-Sitter-like (type II) and anisotropic cosmological solutions (type VI).

Finally, we have shown that the geometries are unconditionally unstable under the radial perturbations. The origin of this instability was shown to be the negativity of the radial pressure ($w_1 < 0$). Our work was focused only on a single matter field which can be responsible both for the asymptotic structure and the black-hole curvature. This result suggests that a cosmological constant (which does not change under perturbation)+matter field model may find a stable solution under the radial perturbations.

Acknowledgment

This work was supported by the National Research Foundation of Korea grants funded by the Korea government NRF-2013R1A1A2006548.

[1] Stephani H., Kramer D., MacCallum M., Hoenselaers C., Herlt E., *Exact Solutions of Einstein's Field Equations,* 2nd ed. Cambridge Monographs on Mathematical Physics, Cambridge University Press, New York (2003).
[2] M. S. R. Delgaty and K. Lake, Comput. Phys. Commun. **115**, 395 (1998) doi:10.1016/S0010-4655(98)00130-1 [gr-qc/9809013].
[3] I. Semiz, Rev. Math. Phys. **23**, 865 (2011) doi:10.1142/S0129055X1100445X [arXiv:0810.0634 [gr-qc]].
[4] Herrera L. and Santos N. O., Phys. Rep. **286**, 53 (1997).
[5] Ruderman R., Ann Rev. Astron. Astrophys. **10**, 427 (1972).
[6] Kippenhahn R. and Weigert A., *Steller Structure and Evolution* Springer, Berlin (1990).
[7] Dev K. and Gleiser M., Gen. Re. Grav. **34** 1793 (2002).
[8] Bhar P, Eur. Phys. J. C **75**, 123 (2015).
[9] B. S. Ratanpal and P. Bhar, arXiv:1612.05417 [gr-qc].
[10] Thirukkanesh S. and Maharaja S. D., Class. Quant. Grav. **25**, 235001-1 (2008).
[11] Ivanov B. V., Phys. Rev. D **65**, 104001-1 (2002).
[12] Varela V., Rahaman F., Ray S., Charkraborty K. and Kalem M., Phys. Rev. D **82**, 044052-1 (2010).
[13] Matese J. J. and Whitman P. G., Phys. Rev. D **11**, 1270 (1980).
[14] Mak M. K. and Harko T., Proc. Roy. Soc. Lond. A **459**, 393 (2003).
[15] Bower R. and Liang E. P., Astrophys. J. **188**, 657 (1974).
[16] I. Cho and H. C. Kim, arXiv:1610.04087 [gr-qc].
[17] M. Barriola and A. Vilenkin, Phys. Rev. Lett. **63**, 341 (1989). doi:10.1103/PhysRevLett.63.341